

CARROT DISEASES

& other factors affecting carrot packout

By Dr. Hoong Pung & Pam Cox, Serve-Ag Research (a division of Serve-Ag Pty Ltd), 16 Hillcrest Road, Devonport, Tasmania 7310, Australia. ABN 36 009 518 674

Corky Crown Rot



Streptomyces species

Corky crown rot does not deteriorate in storage. Usually associated with potatoes as previous crops.

Carrot Scab



Streptomyces species

Scab lesions do not deteriorate storage. Usually associated with potatoes as previous crops.

Tiger Stripe or Ring Rot Disease



Phytophthora species

This disease can develop in the field and sometimes become apparent only after storage or in transit. Usually associated with poor drainage and in ground storage.


Cavity Spot



Pythium sulcatum

The cavities can increase in size in storage. Incidence appears to increase with multiple carrot cropping.

Smooth Crown Rot



Stem infection by *Rhizoctonia* can spread to crown area

Fusarium species,

Rhizoctonia, or *Sclerotinia*.

Usually does not deteriorate in storage. Wet conditions favour this disease, which increases in severity over time.

Sclerotinia Rot



Sclerotinia sclerotiorum

Rot may spread in storage to the rest of the infected carrot, or to adjacent carrots. Wet and warm conditions favour this disease.


Sour Rot



Geotrichum species

This disease can develop in the field and in storage. Associated with poor drainage.

Tip Rot



Tip rot is caused by poor drainage in the field and small cracks that develop on root tips. This usually becomes apparent only during storage.

Black Ring Rot



Black ring rot develops when stem decay, due to bacterial or fungal rot, spreads into the crown tissues. Severe rot can develop if the crop is left in the field long after plant maturity.

Violet Root Rot



Rhizoctonia crocorum

Lesions will enlarge and merge as the disease progresses in the field causing large areas of decay. Shallow lesions on carrots at harvest can enlarge and deteriorate in storage. Associated with poor drainage.

Carrot Black Scurf




Black sclerotia of *Rhizoctonia* on carrot surface.

Rhizoctonia solani

Not normally considered a problem, as the sclerotes are removed during the washing process. The carrots do not deteriorate in storage. Usually associated with potatoes as previous crops.

Carrot Forking



Forking of carrots due to tip damage by *Pythium*.



Forking of carrots due to tip damage by root-knot nematodes.

Factors that can damage root tips, e.g. *Pythium*, root-knot nematodes and compaction, can cause carrot forking.

Carrots are also rejected due to other non-disease factors



Except for the carrots with 'Shadow' and 'Weather Damage' (far left), the non-disease photographs have been reproduced courtesy of Field Fresh Australia.

The Horticultural Research and Development Corporation provided funding for this work, as a part of project VG96015 – "An investigation of carrot diseases in north-west Tasmania and their control." No part of this poster may be copied or reproduced without prior written permission from Serve-Ag Pty Ltd.

